

Placental Delivery The 3rd Stage of Labour

Patient Information

Maternity Service



The Patient Information Leaflets page on the Trust website is available on the link: https://www.wwl.nhs.uk/patient-information-leaflets or scan the QR code.

Author ID: LJ Leaflet ref: Obs 051

Version: 7

Leaflet title: Placental Delivery - The 3rd Stage of Labour

Last review: August 2025 Expiry date: August 2027











Introduction

Your third stage of labour begins once your baby is born and ends when you deliver the placenta (afterbirth). You can choose to have either a physiological (or natural) third stage or an actively managed third stage.

A **physiological 3rd stage** means that you wait for the placenta to be delivered naturally. After your baby's birth, your midwife will delay placing a clamp on the umbilical cord until it has stopped pulsating to allow oxygenated blood to pulse from the placenta to your baby.

Your uterus (womb) will contract, and the placenta will peel away from the wall of your uterus. The placenta will then drop down into your vagina, ready for you to push it out. Breastfeeding and skin-to-skin contact with your baby stimulates contractions and may help your natural third stage along. It may be easier to push your placenta out if you are in an upright position.

Your baby's cord is usually clamped and cut after the placenta has been delivered, or when the cord has stopped pulsating.

Your midwife will check the placenta and membranes to ensure they are complete and that no part has been left behind.

In an **actively managed 3rd stage** your midwife will give you an injection of 1ml Syntometrine in your thigh once your baby's shoulder emerges. After your baby is born cord clamping is delayed for at least one minute and up to five minutes to allow oxygenated blood to pulse from the placenta to your baby. The cord will be clamped and then cut. Your midwife may advise you to have an actively managed third stage if you had complications in your pregnancy or birth.

The injection causes your uterus to contract strongly so the placenta comes away. Once your uterus contracts and there are signs that the placenta is ready to deliver, your midwife will pull gently on the cord while pressing on your tummy above your pelvis to support your uterus while she eases out the placenta.

Both methods take a little over 10 minutes from the birth to when you deliver the placenta. A physiological 3rd stage tends to take longer.

The main advantage of an actively managed 3rd stage is possibly a lower risk of heavy bleeding immediately after the birth. You will also be less likely to become anaemic in the days afterwards. The injection, which is used to narrow your blood vessels and stimulate contractions may occasionally have side effects such as headaches, nausea and a greater need for pain relief.

Your midwife or doctor will recommend you have an actively managed third stage if you had complications during pregnancy or labour, such as:

- a twin pregnancy
- polyhydramnios (too much liquor or water around the baby) or other condition which over-stretches your uterus
- heavy bleeding during pregnancy or in labour
- a low-lying placenta
- anaemia
- an induced or very long labour
- an assisted birth
- a retained placenta
- heavy bleeding after a previous birth

Whilst a physiological 3rd stage is not recommended at caesarean section, it is possible to delay cord clamping for 1 to 5 minutes, providing that your baby does not require immediate resuscitation and you are not bleeding heavily.

Changing from a physiological to an actively managed 3rd stage may be required if you start to bleed more heavily. Your midwife or doctor will treat you for a retained placenta if your physiological 3rd stage takes longer than one hour or your actively managed 3rd stage takes longer than half an hour.

Talk to your midwife about your options for the 3rd stage of labour when you write your birth plan. That way you can discuss your wishes and together, you and your midwife can make a decision which is right for you and your baby.

	Active	Physiological
Length of third stage	shorter	longer
Nausea and vomiting	100 in 1000 women	50 in 1000 women
Haemorrhage > 1 litre	13 in 1000 women	29 in 1000 women
Blood transfusion	14 in 1000 women	40 in 1000 women

References:

Intrapartum care: care of healthy women and their babies during childbirth. Issued: June 2025 NICE Clinical Guideline, Intrapartum Care NG235.

Intrapartum Care for healthy women and babies. guidance.nice.org.uk/cg190.

Comments, Compliments or Complaints

The Patient Relations/Patient Advice and Liaison Service (PALS) Department provides confidential on the spot advice, information and support to patients, relatives, friends, and carers.

Contact Us

Tel: 01942 822376 (Monday to Friday 9am until 4pm)

The Patient Relations/PALS Manager
Wrightington, Wigan and Leigh Teaching Hospitals NHS Foundation Trust
Royal Albert Edward Infirmary
Wigan Lane
Wigan
WN1 2NN

Ask 3 Questions

Become more involved in decisions about your healthcare. You may be asked to make choices about your treatment. To begin with, try to make sure you get the answers to three key questions:

- 1. What are my options?
- 2. What are the positives and negatives of each option for me?
- 3. How do I get support to help me make a decision that is right for me?



How We Use Your Information

For details on how we collect, use, and store the information we hold about you, please see patient information leaflet, Ref. **Corp 006** How we use your information, this can be found on the Patient Information Leaflets page on the Trust website, see details on the front cover.

This leaflet is also available in audio, large print, Braille, and other languages upon request. For more information, please ask in the department/ward.

© Wrightington, Wigan and Leigh Teaching Hospitals NHS Foundation Trust. All rights reserved. Not to be reproduced in whole or in part without the permission of the copyright owner.

Call 111 first when it's less urgent than 999.



Phone: 0808 802 1212

Text: 81212

www.veteransgateway.org.uk

