

Deep Vein Thrombosis

Patient Information

Venous Thromboembolism (VTE) Unit

Author ID: SG
Leaflet Number: VTE 002
Version: 2
Name of Leaflet: Deep Vein Thrombosis
Date of last review: November 2019
Next Review Date: November 2021



VTE Unit

(Deep Vein Thrombosis)

If you have any questions or worries about any of the following information, please feel free to contact the Unit on:

Direct line: 01942 773331

Or via the hospital switchboard: 01942 244000 and ask for extension 3331

The service is available Monday to Friday 9:00 am to 5:00 pm.

Outside of these hours, please contact the Accident and Emergency Department on 01942 822429 or the Ambulatory Assessment Area on 01942 822983.

Today you have visited the hospital because of swelling and pain in your leg, possibly due to a Deep Vein Thrombosis or more commonly known as a DVT.

We have listed below some information for you, which we hope will help you to understand more clearly the sequence of treatment which is likely to follow on from your investigations.

What is a DVT (Deep Vein Thrombosis)?

A DVT is a blood clot which develops in the larger deep veins of your leg.

How do I know for certain that I have a blood clot?

By having a Duplex/Doppler ultrasound scan, you will know for certain.

What treatment will I receive while the tests are being made?

You will be prescribed a drug called Fragmin or Clexane as soon as investigations begin.

1. Following medical assessment and blood test results you will have a Duplex/Doppler ultrasound scan.

2. You will be given an anticoagulant injection called Fragmin or Clexane treatment which will continue until the Duplex/Doppler ultrasound scan has been performed and immediate results are available.

3. If you require a Duplex ultrasound scan it is performed in the Circulation Lab at the Christopher Home, Monday to Friday, 9:00 am to 5:00 pm. Following the procedure you will attend the VTE unit for the results and treatment if needed.

4. If you require a Doppler ultrasound scan it will be performed in the X-ray Department at Wigan Royal Albert Edward Infirmary. Following the scan you will attend the VTE Unit for the results and treatment if needed.

What is a Duplex/Doppler ultrasound scan?

The scans are non-invasive painless ultrasound tests, which examine the veins in the legs.

The tests are performed to detect a thrombus (clot) in the leg.

There is no special preparation for the test, but access is needed to the area to be scanned.

The test usually lasts for 15 to 30 minutes.

When and where will I have this scan?

You will be given the appointment to attend the X-ray Department or Circulation Lab on a next available appointment basis. It is very important that you keep this appointment as suitable and effective treatment depends on the result.

The date and time for your ultrasound scan

.....

at

If you have any problems keeping this appointment, please contact the X-ray Department on 01942 822409 or the Circulation Lab on 01942 822267 as soon as possible.

What are Fragmin and Clexane?

They are anticoagulant drugs that reduce the risk of a blood clot becoming larger or “breaking off”.

How is Fragmin or Clexane administered?

It is given once a day by injection into the skin on the surface of your stomach or upper thigh.

When do I start this treatment?

The first injection will be given by a nurse in hospital, and then can be administered in a number of different ways e.g. yourself, your nearest clinic or District Nurses.

When do the injections stop?

The injections will stop immediately if the Duplex/Doppler is negative. They will continue if the tests are positive for a DVT.

What is the treatment for a DVT?

You will be given an anticoagulant tablet that is taken by mouth. The decision about medication will be discussed in the VTE Unit dependent upon your general health and blood results.

Further information

- You may continue with your normal daily activities.
- If possible keep mobile, when sitting raise your legs unsupported for a few seconds every hour.
- Try to avoid standing for long periods.
- Avoid any contact sport.
- It is safe for you to be treated at home with a DVT.
- Side effects are very rare.
- If you should have any of the following:
 - Unexpected bleeding

- Unexplained bruising
- Breathlessness
- Chest pain

Please report to the Accident and Emergency Department at the hospital and bring your prescription card with you.

Patient notes

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Comments, Compliments or Complaints

The Patient Relations/Patient Advice and Liaison Service (PALS) Department provides confidential on the spot advice, information and support to patients, relatives, friends and carers.

Contact Us

Tel: 01942 822376 (Monday to Friday 9am to 4pm)

The Patient Relations/PALS Manager
Wrightington, Wigan and Leigh NHS Foundation Trust
Royal Albert Edward Infirmary
Wigan Lane
Wigan WN1 2NN

Ask 3 Questions

Become more involved in decisions about your healthcare. You may be asked to make choices about your treatment. To begin with, try to make sure you get the answers to three key questions:

1. What are my options?
2. What are the pros and cons of each option for me?
3. How do I get support to help me make a decision that is right for me?



How We Use Your Information

For details on how we collect, use and store the information we hold about you, please take a look at our “how we use your information” leaflet which can be found on the Trust website: https://www.wwl.nhs.uk/patient_information/leaflets

This leaflet is also available in audio, large print, braille and other languages upon request. For more information please ask in department/ward.

© Wrightington, Wigan and Leigh NHS Foundation Trust
All rights reserved. Not to be reproduced in whole or in part without the permission of the copyright owner

