

# Reducing the Risk of Venous Thromboembolism (Blood Clot)

## Patient Information

VTE Department

Author ID:	CG
Leaflet Ref:	VTE 004
Version:	7
Leaflet title:	Reducing the Risk of Venous Thromboembolism
Last review:	May 2021
Expiry Date:	May 2023

## Introduction

This information leaflet is to help you to understand the care and treatment of people who are at risk of developing a Venous Thromboembolism (VTE) from surgery or during hospital admission as an inpatient. Your healthcare team will be happy to answer any questions you have.

## Venous Thromboembolism

This is the medical term for a blood clot that forms within a vein. The two commonest sites affected are the legs, when it is called a Deep Vein Thrombosis (DVT), or in the lungs, when it is called a Pulmonary Embolism (PE).

A DVT can be painful and cause significant leg swelling, which can make walking difficult, they can also cause a PE to occur. A PE can cause chest pain, difficulty in breathing and they can be fatal.

## Risk Factors

There is a nationally agreed set of risk-factors which all patients having surgery or being admitted to hospital are assessed for.

Example of some risk factors:

- Undergoing surgery
- You or a member of your family have had a blood clot in the past
- You have cancer
- You are taking oestrogen therapy
- You are overweight (body mass index greater than 30)
- You are unable to move around
- You are over 60 years old

If you have one or more of the risk-factors, then you will be considered for ways to reduce your risk of developing a VTE.

## Reducing the risk of VTE

There are two main ways hospitals have of reducing your risk of developing a VTE:

1. Using compression stockings
2. Using medicine

## Compression Stockings

Compression stockings are tight legwear specially designed to reduce the risk of a DVT. The stockings squeeze your feet, lower legs, and thighs, helping the blood to move around your body better. They are more commonly used for patients requiring surgery, rather than patients admitted to hospital for other reasons.

If you are offered compression stockings it is important to wear them as much as possible, until you are back to your usual level of activity.

## Medicines

Depending on your risk factors, your healthcare professional may also offer you medicine during your hospital stay. The commonest medicine currently used in this Trust to prevent VTEs is called Dalteparin (brand name – Fragmin). It is an injection given once a day into your tummy or thighs. Some patients requiring surgery to their knees or hips may be offered a tablet instead of an injection.

The medicines used to prevent VTEs can increase the risk of bleeding. Therefore, everyone who is considered for one of these medicines will also be assessed for their risk of bleeding according to national guidance. If your risk of bleeding is felt to be higher than your risk of a VTE, your healthcare professional will discuss this with you in more detail.

## Additional ways to reduce your risk

Your healthcare team will encourage you to move about or help you to undergo leg exercises. It is also important to drink plenty of fluids where possible to help reduce your chances of getting a blood clot. If you are struggling to drink, then your healthcare team may give your fluids through a drip into your vein.

## After discharge from hospital

After being discharged from hospital some patients may still be at a high risk of developing a VTE. Therefore, some patients are continued on a medicine or advised to wear stocking for several more weeks. Your healthcare professional will explain this in more detail if this applies to you.

## How to tell if you have a VTE

There are certain signs to look out for that could mean you have a VTE. You should tell your healthcare team or GP (if at home) immediately if you experience any of the following in the days or weeks after your inpatient admission or surgery:

- You have pain or swelling in your leg
- The skin on your leg is hot or discoloured (red, purple or blue)
- Your feet are numb or tingling
- The veins near the surface of your legs appear larger than normal

- You become short of breath
- You feel pain in your chest, back or ribs which gets worse when you breathe deeply
- You cough up blood

For further information please speak to a member of your healthcare team or GP.

## Reference

<https://www.nice.org.uk/guidance/ng89> NICE NG89; March 2018

Please use this space to write notes or reminders.

---

## Comments, Compliments or Complaints

The Patient Relations/Patient Advice and Liaison Service (PALS) Department provides confidential on the spot advice, information and support to patients, relatives, friends and carers.

## Contact Us

Tel: 01942 822376 (Monday to Friday 9am to 4pm)

The Patient Relations/PALS Manager  
Wrightington, Wigan and Leigh Teaching Hospitals NHS Foundation Trust  
Royal Albert Edward Infirmary  
Wigan Lane  
Wigan WN1 2NN

---

## Ask 3 Questions

Become more involved in decisions about your healthcare. You may be asked to make choices about your treatment. To begin with, try to make sure you get the answers to three key questions:

1. What are my options?
2. What are the pros and cons of each option for me?
3. How do I get support to help me make a decision that is right for me?



---

## How We Use Your Information

For details on how we collect, use and store the information we hold about you, please take a look at our “how we use your information” leaflet which can be found on the Trust website: <https://www.wwl.nhs.uk>

---

This leaflet is also available in audio, large print, Braille and other languages upon request. For more information please ask in the department/ward.

---

© Wrightington, Wigan and Leigh Teaching Hospitals NHS Foundation Trust  
All rights reserved. Not to be reproduced in whole or in part without the permission of the copyright owner.

**Call 111 first when it's less urgent than 999.**



Phone: 0808 802 1212  
Text: 81212  
[www.veteransgateway.org.uk](http://www.veteransgateway.org.uk)

