

Bulkamid

Patient Information

Gynaecology Department



The Patient Information Leaflets page on the Trust website is available on the link:
<https://www.wwl.nhs.uk/patient-information-leaflets> or scan the QR code.

Author ID: JD
Leaflet ref: Gyn 050
Version: 7
Leaflet title: Bulkamid
Last review: September 2021
Expiry Date: September 2023



Bulkamid is one of the treatment options for stress incontinence

What is stress incontinence?

The muscles of the pelvic floor support the bladder and usually help keep it closed or open as necessary. Stress incontinence usually happens when these muscles become weak, so when there is sudden extra pressure “stress” on your bladder, it cannot stay closed as it should and some urine leaks out. This leakage happens during normal everyday activities, and most often when you cough, sneeze, laugh, exercise, or change position. Whether you leak a small or large amount of urine, stress incontinence can be embarrassing and distressing.

Stress incontinence can be triggered by pregnancy, childbirth, or the menopause but sometimes there is no clear cause for it.

Do I need treatment?

What you do about stress incontinence will depend on how far it affects you and what you feel you can cope with as well as your general health.

What treatments are available?

Practical advice from a continence nurse specialist on managing your daily life such as:

- Adjusting your daily routines to help you cope better
- How you can help yourself by losing weight if you are overweight
- Managing a chronic cough if you have one
- Giving up smoking

Physiotherapy

Special physiotherapy exercises to make your pelvic floor muscles stronger and improve control of your bladder. These carry up to 70% chance of improving or curing symptoms. It is usually recommended that a programme of physiotherapy be undertaken prior to considering any surgery.

Alternatives

Tablets

A tablet called Duloxetine (Yentreve) can help stress incontinence for some people though it only works whilst you are taking it and does not suit everyone.

Surgery

Not everyone with stress incontinence needs an operation. Whether you choose to have surgery will depend on how far stress incontinence affects your daily life and what you feel you can cope with. You may want to consider surgical options if other things (such as exercises to help strengthen the muscles in the pelvic floor) have not helped.

Surgical procedures for stress incontinence are not usually suitable if you still plan to have children or think you might want to in the future. Before any surgical procedure you will be asked to attend for tests on your bladder (urodynamic studies) to confirm that the problem is stress incontinence and therefore likely to be helped by surgery.

Surgical options available in Wigan and Leigh are:

- Colposuspension
- Fascial sling
- Bulkamid (urethral bulking agent)

A colposuspension is a fairly major operation which can often be carried out using key-hole surgery and has an 80-85% success rate.

A fascial sling is a major abdominal operation which uses tissue taken from your abdomen to make a sling which can be inserted to support your bladder. It has similar success rates to a colposuspension.

Bulkamid has a lower cure rate than other procedures but there are fewer complications, and it does not require an anaesthetic. It can be performed in an outpatient clinic with a recovery time of only one to two days and is suitable even if you plan further children.

Each procedure is described in its own leaflet.

Bulkamid

Bulking agents are a relatively non-invasive surgical treatment option, using local anaesthetic, for stress incontinence and an alternative to major surgery. Bulking agents are injected into the wall of the urethra (tube passing from your bladder to the outside through which you pass urine) to improve the sealing mechanism of the bladder. The treatment takes about 15 to 20 minutes and involves no hospital stay or lengthy recovery period.

Bulking agents can be offered to women who are reluctant or not medically fit enough to have more major surgery or who are planning to have more children. This does not rule out the possibility of another surgical treatment at a later date.

The bulking agent used in this Trust is Bulkamid. Unlike some other bulking agents this is permanent. Enzymes within the body do not break it down and it is not reabsorbed by the body. It is soft and does not induce scarring around it and no allergies have been reported. It has been used in plastic surgery and to make contact lenses for over 10 years.

The procedure is carried out in the outpatient department at Leigh Infirmary. Some local anaesthetic jelly is squeezed into the urethra (tube from bladder to outside) 5 minutes

before the procedure is commenced. Two local anaesthetic injections are given around the area. A small tube with a camera attached (cystoscope) is placed into the urethra and some fluid is put into the bladder. Four injections of Bulkamid are placed into the urethral wall via this cystoscope. The procedure takes 15 to 20 minutes.

The procedure may be uncomfortable, and your bladder may feel rather full. Women sometimes describe a sensation of stinging as the Bulkamid is injected but this wears off soon afterwards.

Following the procedure you will wait in clinic until you pass urine. If your bladder does not feel fully empty after you have passed urine your bladder will be scanned to check it has emptied properly and then you will be able to go home. As the local anaesthetic can make you feel a little wobbly after the procedure, we recommend that someone comes with you and that you avoid driving until the following day.

Sometimes a single treatment is not sufficient and one further treatment some months later is needed.

If you have managed to empty your bladder in clinic it is very rare for you to have any problems later but if you, do it is important to seek help before you become too uncomfortable.

Please phone Swinley Ward on 01942 822016 to let them know you need to come up and then attend the Gynaecology emergency room on Swinley Ward, Wigan Infirmary.

Benefits

Overall it is reported to cure or substantially improve stress incontinence in 60-70% of women. There are not yet any long term studies of success but studies of women one to two years after the procedure suggest that the results are very stable over that time.

It is not designed to help urinary frequency, urgency, or leakage due to urgency however, they may improve.

Complications

There are very few reported complications.

- You may experience discomfort and pain while passing urine after the procedure. This should resolve after 24 hours and can be managed with mild pain killers.
- A urine infection may occur, but you will be given a short course of antibiotics to attempt to prevent this.
- There might be a small amount of blood staining of the urine for a day or two.
- Difficulty in passing urine. Rarely you may experience difficulty in passing urine after the procedure. This usually resolves within the first 24 hours and your bladder may need to be emptied with a catheter during that time. Very rarely, you may need to go

home with a catheter for a few days. You will then have to return to the ward for the catheter to be removed and to make sure that your bladder is emptying properly.

- The procedure may fail to completely cure your stress incontinence but, in this case, there is an option to repeat the procedure.

Recovery

There should be minimal discomfort after the procedure and a full recovery would be expected within one to three days. There are no limitations on activities following it. Improvement in urinary leakage would be expected almost immediately.

Contact details

If you have any questions or concerns, please contact:

Hanover Women's Healthcare Unit telephone number 01942 264962 or 01942 264963
(Monday to Friday 9:00am to 4:30pm)

Swinley Ward on 01942 822016

Comments, Compliments or Complaints

The Patient Relations/Patient Advice and Liaison Service (PALS) Department provides confidential on the spot advice, information and support to patients, relatives, friends, and carers.

Contact Us

Tel: 01942 822376 (Monday to Friday 9am to 4pm)

The Patient Relations/PALS Manager
Wrightington, Wigan and Leigh Teaching Hospitals NHS Foundation Trust
Royal Albert Edward Infirmary
Wigan Lane
Wigan WN1 2NN

Ask 3 Questions

Become more involved in decisions about your healthcare. You may be asked to make choices about your treatment. To begin with, try to make sure you get the answers to three key questions:

1. What are my options?
2. What are the pros and cons of each option for me?
3. How do I get support to help me make a decision that is right for me?



How We Use Your Information

For details on how we collect, use, and store the information we hold about you, please see patient information leaflet, Ref. **Corp 006** How we use your information, this can be found on the Patient Information Leaflets page on the Trust website, see details on the front cover.

This leaflet is also available in audio, large print, Braille, and other languages upon request. For more information, please ask in the department/ward.

© Wrightington, Wigan and Leigh Teaching Hospitals NHS Foundation Trust.
All rights reserved. Not to be reproduced in whole or in part without the permission of the copyright owner.

Call 111 first when it's less urgent than 999.



EMPLOYER RECOGNITION SCHEME

SILVER AWARD

Proudly supporting those who serve.

Phone: 0808 802 1212
Text: 81212
www.veteransgateway.org.uk

