



**Wrightington, Wigan and
Leigh Teaching Hospitals**
NHS Foundation Trust

Ward 1 - Discharge Advice

Ward One - Discharge Advice

Patient, Relative and Carer Information

Wrightington Hospital

- Author ID: IJ
- Leaflet ref: Musc 034
- Version: 8
- Leaflet title: Ward One - Discharge Advice
- Last review: May 2025
- Expiry date: May 2027

Introduction

Thank you for attending Ward One, we hope you have found everything to your satisfaction. If you do have any concerns, please raise them with a member of staff so we can deal with them promptly.

The following booklet will provide you with the relevant information required for your ongoing care needs when you have gone home. This is also intended to provide advice to patients' relatives and / or carers to support our patients on their discharge from hospital.

Please do not hesitate to contact us at any time by telephoning:

Ward One Monday to Friday, 7:30am until 7:30pm

Male Bay 01257 256550

Female Bay 01257 256551

Outpatients Department: 01257 256299, Monday to Friday 9am until 5pm

Outpatients

Your clinic appointment is on:

At: Royal Albert Edward Infirmary Wrightington Leigh

(Staff to tick as appropriate)

If your appointment is not given on discharge, you should receive it within 1 week. If for any reason you do not receive an appointment through the post, or if you have any problem with your outpatient appointment, please contact:

01257 256222 or 01257 256241

Wound care

- Care needs to be taken to keep dressings clean and dry. Please ask for advice regarding bathing precautions prior to your discharge.
- Your wound should be left alone YES/NO
- Where applicable, the outer bandaging needs to be removed down to the wound dressing in days.
- Wound dressings should be changed as advised by Nursing Staff on Discharge.
- If applicable, your stitches should be removed in days. This should be carried out at:

Wrightington,
Wigan or Leigh
Outpatients

District
Nurse

Nur
Out
Clin

Other.....

(Staff delete as appropriate)

You have been referred to Therapy. YES NO

How to change your dressing

- Only change your dressings if necessary.
- Ensure you clean your hands prior to carrying out the procedure. Use soap and water and dry with a clean towel.
- Remove outer dressing. If adhesive dressing is clean and not blood stained, leave intact.
- If not, carefully remove adhesive dressing, ensuring you do not touch the wound.
- Apply the dressing supplied. This is done as follows: Remove the dressing from the outer packaging. Peel off cover underneath and smooth the dressing over the wound. Please ensure you do not touch the white central pad. Then carefully peel off the top cover to leave the dressing in place. Dressings are shower proof.
- If, at any point during the procedure you are concerned, please contact the ward or your General Practitioner (GP).

Swelling

It is a natural part of the healing process for localised swelling to occur after surgery. To reduce swelling, your limb needs to be elevated. Use additional pillows / cushions to rest it up at night or whilst sitting during the day.

If you have had any shoulder surgery, please commence exercises, and use the sling as per physiotherapy instructions. It is normal for shoulder arthroscopy procedures to ooze for the first couple of days.

Should you become concerned about any of the following:

- Any colour changes.
- Increased redness, with hot / burning pain and raised temperature to the operated area.
- Further numbness / pins and needles.
- Swelling that has increased beyond the size of the dressing / splint which has become painful.
- The dressing / splint has become too loose or too tight.

Do not hesitate to contact the ward or Therapy Department for advice.

If you have any problems regarding therapy, please contact:

Upper and Lower Limb Physiotherapy: 01257 256307, 8.30am until 5pm

Hand Unit Office: 01257 488272, 8.30am until 5pm

Pain Control

Pain is a normal experience after surgery. It is important to actively help your body to recover from surgery by following any instructions given by the hospital.

If you have any worries about your pain at home, please seek advice from a medical professional.

Driving

Please discuss this with your Consultant, who will provide advice regarding when it is safe to drive following your surgery.



Version number: **8**
Last modified date: **13th June 2026**

All rights reserved © 2026
WWL Teaching Hospitals NHS Foundation Trust