Wrightington, Wigan and Leigh Teaching Hospitals
NHS Foundation Trust

Ward 3 Eye Unit - Injection Clinic

Version number: **2**
Last modified date: **12th November 2026**

Ward 3 Eye Unit – Injection Clinic

Patient Information

Ophthalmology Services

- Author ID: CR
- Leaflet ref: Ophth 023
- Version: 2
- Leaflet title: Ward 3 Eye Unit - Injection Clinic
- Date Produced: March 2025
- Expiry Date: March 2027

What are the risks with these injections?

As with any medical procedure, there is a small risk of complications that may occur following an intravitreal injection. Most complications that might occur are from the injection itself, rather than the drug.

For most patients, the benefit of the treatment outweighs the small risk of injection injury. This should have been discussed with you when you started your course of treatment.

The following are the main potential risks and side effects of anti-VEGF eye drug injections, but this is not a complete list of all risks. These risks are all rare. Significant loss of vision due to this treatment is very uncommon.

- You might notice some discomfort and redness and your vision may be blurred for the first few days after your treatment and this usually settles between 24 – 48 hours.
- You may notice floaters in your eye which usually disappear after a few days.
- Sometimes the eye may feel gritty and watery. This should settle down after a day.
- It is recommended that you take your normal analgesia after the injection if you eye feels uncomfortable.
- Serious eye infection - If your eye becomes progressively red, sensitive to light, swollen with increased pain, or if your vision gets worse after the injection treatment, you must seek medical help and advice from an ophthalmologist.

This might indicate infection and normally occurs within the first few days after the injection. The risk is minimal, but this is the most important complication, and you must contact us immediately if you are concerned about this.

Important Information: Your vision may be at risk if you do not complete the course of treatment. Boston House will arrange your next appointment for you.

How to contact us:

Please contact Boston House if you have any concerns, the unit is open from 08:30am – 5pm Monday to Friday.

The contact number is **01942 773116** or **01942 822244**.

Outside of these hours you may attend your local A&E department.

A&E department at RAEI Wigan is open 24 hours a day and is contactable on 01942 822440.

They may refer you to a specialist Eye Unit at Manchester or Liverpool. If you wish to contact Manchester or Liverpool Eye Units directly when Ward 3 Eye Unit is closed, they are available on the following numbers:

Manchester Royal Eye Hospital:

0161 2765597

Royal Liverpool Eye Hospital:

0151 706 2000

Why an eye injection may be cancelled on the day:

- If you are unwell, e.g. urine, chest or eye infection, heavy cold, eye or are taking antibiotics for any infection.
- If you have recently had a heart attack or a stroke.
- Are pregnant or breast feeding.
- If you have had any vaccinations in the last 7 days (vaccinations should be avoided for 1 week either side of your injection).

Please leave contact lenses out for 3 days prior & post injection.

If any of the above apply, please contact: 01942 822244, Monday-Friday 8:30am to 5pm.

Leigh Infirmary

The Avenue
Leigh
Lancashire
WN7 1HS

Main Phone: 01942 244 000
Walk-in-Centre: 01942 483 453



Version number: **2**
Last modified date: **12th November 2026**

All rights reserved © 2026
WWL Teaching Hospitals NHS Foundation Trust