



**Wrightington, Wigan and
Leigh Teaching Hospitals**
NHS Foundation Trust

Prostap and Zoladex Therapy

Prostap and Zoladex Therapy

Patient Information

Gynaecology Department

- Author ID: UG
- Leaflet Ref: Gyn 043
- Version: 10
- Leaflet title: Prostap and Zoladex Therapy
- Last review: April 2025
- Expiry Date: April 2027

Introduction

Prostap and Zoladex are types of drugs called gonadotrophin-releasing hormone analogues (GnRHa). They are very similar; Prostap is usually given in our hospital. In gynaecology they may be used to treat fibroids, endometriosis, premenstrual syndrome (PMT) and for In Vitro Fertilisation (IVF) treatment. They are also used in the treatment of various conditions including prostate and breast cancer.

Possible side effects

Prostap and Zoladex work by causing a temporary 'artificial menopause' in the body, by decreasing the production of oestrogen and by stopping menstruation (periods). Some women experience increased bleeding in the first month on treatment although this usually resolves by itself. The main side effects of the drugs are similar to the effects of the menopause:

- Hot flushes
- Increased sweating
- Vaginal dryness causing discomfort during intercourse
- Decreased libido (sex drive)
- Decreased bone density (thinning of the bones) if used for a long period of time without HRT (see below)

In order to minimise these side effects patients are also prescribed a form of Hormone Replacement Therapy (HRT) called Livial, which is taken in the tablet form every day. The Royal College of Obstetricians and Gynaecologists has stated that the safe length of time

that GnRHa can be used is still unclear, but treatment as long as it is combined with HRT appears to be safe and effective for up to two years. As far as can be determined, the treatment with HRT does not carry any risks in those under 50 years old.

If you experience any other side effects, you should discuss these with your doctor or nurse.

Alternatives

There may be alternative treatments for the condition for which Prostag is being offered. Tablet form of GnRH analogue is now available called Ryeqo, this tablet has HRT in it. Therefore, separate HRT (Livial) is not required. If you would like to have more information about this, please consult the doctor treating you. These alternatives will vary depending on the condition being treated.

Bone density scans

Some people stay on treatment for longer than two years. If this happens you would then be asked to have a DEXA scan, (similar to an x-ray), to measure your bone density. If you stay on treatment for longer than this, you will need to have a bone density scan every two years.

How to take Prostag and Zoladex

It is given as an injection during the first five days of your period and then given every month or every three months depending on which suits you best.

Barrier methods of contraception should still be used whilst on Prostag or Zoladex.

Contact information

If you have any questions, please contact:

Hanover Women's Unit (9am until 5pm Monday to Friday) telephone **01942 264959**

Swinley Ward (24 hours) telephone **01942 822568**



Version number: **10**
Last modified date: **03rd July 2026**

All rights reserved © 2026
WWL Teaching Hospitals NHS Foundation Trust