

Simple wrist fracture

Simple Wrist Fracture

Patient Information

Trauma and Orthopaedics

- Author ID: VL
- Leaflet Ref: Musc 083
- Version: 1
- Leaflet title: Simple wrist fracture
- Date Produced: April 2024
- Expiry Date: April 2026

Introduction



You have broken (fractured) a bone in your wrist. This injury is very common. Pain, swelling and bruising to the wrist, hand and fingers is normal and can last for a few weeks.

Because your bones are still in a good position, this injury does not need surgery.

You should make a very good recovery if you follow the advice and do the exercises little and often.

Injury recovery

- It takes 6-12 weeks to heal

- Pain and swelling are normal
- Take simple painkillers e.g. paracetamol for as long as you need to; do not take more than the recommended dose. Your local Pharmacist is a useful source for pain relief information
- Attempt to make a full fist and straighten your fingers as early as possible
- You can start doing gentle wrist exercises out of the splint after 4 weeks
- Smoking slows down healing. We advise you to stop. Please talk to your General Practitioner (GP) or go online at <https://www.nhs.uk/smokefree>

Occasionally a plaster cast is applied instead of a splint. The Fracture Clinic will advise when this should be removed.

Caring for your injury

Elevate your hand for the first 24 – 72 hours after your injury. Sit more than usual and keep your hand lifted on a cushion (chest high). This will help to reduce the swelling.

Using ice can help with your pain and swelling. Wrap a damp tea towel around a bag of frozen peas and place on your wrist and hand for up to 15 minutes, each hour, as needed. Do not use ice if the feeling in your hand is reduced, or if you have skin problems.

Remember to wear your splint for the first four weeks. You can take the splint off to wash and when resting, if comfortable.


Exercise week 1- 4

Gentle movement of the elbow, fingers and thumb creates blood flow to the injury. This helps it to heal and reduces swelling. Starting early prevents stiffness.



Exercise from 4 weeks

Remove the splint and start wrist exercises as below. All exercises should be performed little and often or at least four times per day, but you should avoid pushing into any sharp pain. You can access more information in our [‘removal of cast following a hand or wrist fracture’](#) leaflet TPY 043 (click link or visit wwl.nhs.uk website)



Disclaimer

Not all exercises are suitable for everyone, and while every precaution has been taken in selecting these exercises, they do not come without risks. If you have any concerns regarding your injury, or whether you should partake in the exercises, then you should consult your General Practitioner (GP) or Physiotherapist before beginning any exercises in this leaflet.

It is quite normal to feel some discomfort following exercise, but it is important that this discomfort eases within 2 hours of finishing the exercise and does not have an overall negative effect on your symptoms.

The exercise plans shown in this leaflet are designed to be used as a guide and a point of reference; they can be modified and progressed under the supervision of your Physiotherapist.

Osteoporosis?

If you have broken a bone AND are over 50 years of age, you may be at risk of having osteoporosis (weakening of the bones). You will be contacted separately by the Specialist Fracture Liaison Team and asked to opt in for a review of your bone health. If you live outside the Wigan area, please speak to your GP.

As we get older, bones do not renew themselves as well. This causes them to weaken; it happens to everyone to some degree, but if the bones become fragile, it's called osteoporosis. You can help with certain lifestyle factors:

- Get plenty of calcium and Vitamin D as part of a well-balanced diet and outdoor activities
- Regular weight bearing exercise e.g., walking, running
- Stop smoking
- Do not drink too much alcohol

Frequently asked questions

Driving

You must not drive with the splint / cast on. It is your own responsibility. You are allowed to drive when:

- You can safely control the car
- You are no longer using the splint / cast
- You can grip the steering wheel / gear stick safely
- You can perform an emergency stop

Work

This depends on your individual situation. You can return to work when you feel able to do your job. Consider a phased return where needed / possible.

Sports

You should avoid contact / ball sports for 8-12 weeks. You can exercise your lower body as soon as you feel able and safe to.

For more information online, please visit <https://www.wvl.nhs.uk/virtual-fracture-clinic>

Remember....

- Wrist fractures can be very painful
- The majority of injuries will recover in 6-12 weeks, but you may experience stiffness for up to 12 months
- Gentle exercises will aid your recovery and reduce risk of re-injury

Further Advice

If you have any concerns regarding your wrist, contact:

Fracture Clinic Helpline 01942 822595

Please leave a message with name, telephone number and brief description of reason for call. We will aim to call you back within 24 hours Monday to Friday 8.30am until 5:00pm (please note it will be Monday if you call over the weekend).

If you have any urgent concerns out of hours, please call 111.

Adult MSK Physiotherapy Self-Referral

If you are struggling to regain your movement or get back to normal activities, you can self-refer to Physiotherapy. A referral form can be found on this webpage:

<https://www.wwl.nhs.uk/adult-msk-physiotherapy-self-referral>



Please scan the QR Code to access the website.

Telephone Numbers:

Boston House Health Centre Telephone 03007071113

Leigh Infirmary Telephone 03007071597 / 03007071595

Platt Bridge Health Centre Telephone 03007071772



Version number: 1
Last modified date: **03rd July 2026**

All rights reserved © 2026
WWL Teaching Hospitals NHS Foundation Trust