



**Wrightington, Wigan and  
Leigh Teaching Hospitals**  
NHS Foundation Trust

# Volar plate sprain

# Volar Plate Sprain

## Patient Information

## Trauma and Orthopaedics

- Author ID: VL
- Leaflet Ref: Musc 088
- Version: 1
- Leaflet title: Volar plate sprain
- Date Produced: April 2024
- Expiry Date: April 2026

## Introduction



The volar plate is a thick ligament which joins two bones in the finger. This injury occurs when the finger is bent too far the wrong way, spraining, or tearing the ligament. This can often happen playing sports.

Symptoms include pain, bruising and swelling. An X-ray of your hand may show a very small piece of bone; this happens where the ligament is attached and gets stretched suddenly during the injury. This injury is due to a sprain of the ligament, not a broken bone.

## Injury recovery



- It takes 6-12 weeks to recover from this injury. Often this is much quicker in children
- Pain and swelling are normal
- Wear strapping for up to two weeks as tolerated
- Take simple painkillers e.g., paracetamol for as long as you need to; do not take more than the recommended dose. Your local Pharmacist is a useful source for pain relief information
- A splint can be worn for up to two weeks (if provided); not everyone needs this

**It is very important to start your exercises as soon as possible.**

Smoking slows down the healing process. We advise that you stop (at least whilst recovering from your injury). Speak to your General Practitioner (GP) or go online at: <https://www.nhs.uk/smokefree>

## Exercise



It is very important to maintain movement of the finger to avoid stiffness from DAY ONE, even if this means overcoming some discomfort. Do these exercises every hour during the day (you can do them with tape on). Take splints off to exercise (if provided).

1. Try to bend your fingers, aiming to touch your palm with finger tip.
2. Lay your hand on a flat surface to fully straighten your finger.

It is quite normal to feel some discomfort following exercise, but it is important that this discomfort eases after finishing the exercise and does not have an overall negative effect on y

our symptoms.

## Warning signs!

**Some people as they heal can develop a bend in the finger called a 'contracture'. Contractures can be stubborn, if not treated early on. If you are worried or have NOT got full range of movement by two weeks you should call the Fracture Clinic helpline to arrange an appointment with our Hand Therapy Team.**

It is common for some swelling to stay for many months, and sometimes this can remain forever, but this should not affect the ability to use your hand after six weeks.

Most injuries heal well without any problems; however it may take 3-6 months for your symptoms to settle completely. If you have any worries, please call the helpline.

## Disclaimer

Not all exercises are suitable for everyone, and while every precaution has been taken in selecting these exercises, they do not come without risks. If you have any concerns regarding your injury, or whether you should partake in the exercises, then you should consult your GP or Physiotherapist before beginning any exercises in this leaflet.

The exercise plans shown in this leaflet are designed to be used as a guide and a point of reference; they can be modified and progressed under the supervision of your Physiotherapist.

For more information online please visit <https://www.wvl.nhs.uk/virtual-fracture-clinic>

## Frequently asked questions

---

### When can I drive?

---

It is your own responsibility. You are allowed to drive when:

- You can safely control the car
- You can grip the steering wheel or gear stick
- You can perform an emergency stop

### When can I go back to work?

---

This depends on your individual situation. You can return to work when you feel able to do your job.

## Further Advice

If you have any concerns regarding your finger, contact:

**Fracture Clinic Helpline** 01942 822595 please leave a message with name, telephone number and brief description of reason for call. We will aim to call you back within 24 hours Monday to Friday 8.30am until 5:00pm (please note it will be Monday if you call over the weekend).

## Adult MSK Physiotherapy Self-Referral

If you are struggling to regain your movement or get back to normal activities, you can self-refer to **Physiotherapy**. A referral form can be found on this webpage:

<https://www.wvl.nhs.uk/adult-msk-physiotherapy-self-referral>



Please scan the QR Code to access the website.

### Telephone Numbers:

**Boston House Health Centre** Telephone 03007071113

**Leigh Infirmary** Telephone 03007071597 / 03007071595

**Platt Bridge Health Centre Telephone 03007071772**



Version number: 1  
Last modified date: **03rd July 2026**

All rights reserved © 2026  
WWL Teaching Hospitals NHS Foundation Trust