



**Wrightington, Wigan and
Leigh Teaching Hospitals**
NHS Foundation Trust

Discharge Advice following Cardiac Angiography - Radial

Discharge Advice following Cardiac Angiography (Radial)

Patient Information

Cardiac Catheter Laboratory

- Author ID: CW
- Leaflet ref: CCL 001
- Version: 7
- Leaflet title: Discharge Advice following Cardiac Angiography (Radial)
- Last review: August 2024
- Expiry date: August 2026

Discharge Advice following Cardiac Angiography (Radial)

- You should be accompanied home and have a friend/relative to stay with you over night.
- If severe bleeding occurs at the puncture site, apply firm pressure, and raise your arm. Ask your friend/relative to ring 999 and return immediately to your nearest Accident & Emergency Department.
- If you experience fever, bleeding, persistent tenderness or swelling, redness or warm to touch, numbness or pain at the site, discharge, or rash you must go immediately to your nearest A&E department.
- Modify activities for 48 hours – no straining, lifting greater than 5kg or weightbearing in the arm used to access your arteries.
- It is advisable to drink plenty of fluids, especially water over the next few days; this will assist with flushing the dye out of your body.
- It is advisable to refrain from the consumption of alcohol for 24 hours.
- Do not have a bath or shower until the following day and take extra care when washing and drying the puncture site.
- Refrain from driving for at least 48 hours following your angiogram and 7 days following an angioplasty.
- If you take Metformin **do not** take for 48 hours following your angiogram.

- If you take, Warfarin, Apixaban, Dabigatran, Edoxaban or Rivaroxaban you may restart it on the evening following your angiogram unless the Doctor has instructed otherwise.

Please contact the Cardiac Catheter Lab on 01942 773392 if you have any questions.



Version number: **7**
Last modified date: **13th June 2026**

All rights reserved © 2026
WWL Teaching Hospitals NHS Foundation Trust